

## INFLAMMATORY BOWEL DISEASE SPECIALTY CARE PROGRAM

Phone: **844-805-3357** • Fax: **888-978-6171** 



	T INFORMAT	ION:			CRIBER INFORM		any care neu	
Address:				_ Address: _				
City:		State: Zip: _		City:		State: Z	Zip:	
		Alt. Phone:						
		M OF Caregiver:						
Height:	Weight:	Allergies:		Office Contact:		Phone:		
ate of Diagnosis:			Serious or active infection present?  Yes No  Hep B ruled out or treatment started?  Yes No  TB Test: Positive Negative Date:		dical Documentation)  Prior Failed Treatments:  5-ASA Biologics	Indicate Drug Name and Length of Treatment:		
					☐ Corticosteroids ☐ Immunosuppressant ☐ Methotrexate	s		
Date of Diagnosis: Diagnosis: □ Irritable Bowel Syndrome □ 564.1 Irritable Bowel Syndrome					□ Surgery □ Other			
_		<b>DRMATION:</b> (Please	be sure to	choose both		nance dose wh		
Medicatio		Dosage & Strength			Direction			Refills
☐ Prefilled Syringe Starter ☐ CIMZIA® ☐ 200mg/ml Prefilled Syrin ☐ 200mg Lyophilized Pow		ige 🗆 N	☐ Induction Dose: Inject 400mg SC on day 1, 14 and 28☐ Maintenance: Inject 400mg SC every 4 weeks☐			2	0	
□ HUMIRA		Crohn's Disease/ Ulcerative Colitis Starter			ction Dose: Inject 160mg SC on day 1, then g SC on day 15, then switch to maintenance dose		6	0
		l 40mg/0.8ml Pen l 40mg/0.8ml Prefilled Syl	yringe		nject 40mg SC every other week		2	
□ SIMPO	□ SIMPONI® □ 100mg/ml Smartject® Autoinjector			☐ Induction Dose: Inject 200mg SC at week 0, 100mg SC at week 2 and then switch to maintenance dose			3	0
		□ 100mg/ml Prefilled Syringe □ M			Maintenance: Inject 100mg SC every 4 weeks			
☐ UCERI	ERIS <sup>®</sup> □ 9mg Tablets		ПΤ	☐ Take one tablet daily in the morning with or without food			30	1
☐ XIFAXA	XIFAXAN® □ 550mg Tablets		ПΤ	☐ Take one tablet three times daily for 14 days			42	
┃ □								
5 INJECT	ION TRAININ	IG: O Pharmacist to P	rovide Train	ing O Patient	t Trained in MD Office	O Manufacture	r Nurse	Support
		Y: O Patient's Home  WATION: Please Inclu						